



HAL
open science

Dynamic Arterial Elastance Is Associated With the Vascular Waterfall in Patients Treated With Norepinephrine: An Observational Study

Stephane Bar, Maxime Nguyen, Osama Abou-Arab, Herve Dupont, Belaid Bouhemad, Pierre-Gregoire Guinot

► To cite this version:

Stephane Bar, Maxime Nguyen, Osama Abou-Arab, Herve Dupont, Belaid Bouhemad, et al.. Dynamic Arterial Elastance Is Associated With the Vascular Waterfall in Patients Treated With Norepinephrine: An Observational Study. *Frontiers in Physiology*, 2021, 12, pp.583370. 10.3389/fphys.2021.583370 . hal-03575639

HAL Id: hal-03575639

<https://u-picardie.hal.science/hal-03575639>

Submitted on 12 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Dynamic Arterial Elastance Is Associated With the Vascular Waterfall in Patients Treated With Norepinephrine: An Observational Study

Stéphane Bar^{1*}, Maxime Nguyen^{2,3}, Osama Abou-Arab¹, Hervé Dupont¹, Belaid Bouhemad² and Pierre-Grégoire Guinot^{2,3}

OPEN ACCESS

Edited by:

Lacolley Patrick,
Institut National de la Santé et de la
Recherche Médicale (INSERM),
France

Reviewed by:

Dimitrios Terentes-Printzios,
University of Oxford,
United Kingdom
Thomas Desaive,
University of Liège, Belgium

*Correspondence:

Stéphane Bar
stephane.bar.sb@gmail.com

Specialty section:

This article was submitted to
Vascular Physiology,
a section of the journal
Frontiers in Physiology

Received: 14 July 2020

Accepted: 24 March 2021

Published: 04 May 2021

Citation:

Bar S, Nguyen M, Abou-Arab O,
Dupont H, Bouhemad B and
Guinot P-G (2021) Dynamic Arterial
Elastance Is Associated With the
Vascular Waterfall in Patients Treated
With Norepinephrine: An
Observational Study.
Front. Physiol. 12:583370.
doi: 10.3389/fphys.2021.583370

¹Department of Anaesthesiology and Critical Care, Amiens University Hospital, Amiens, France, ²Department of Anaesthesiology and Critical Care, Centre Hospitalier Régional Universitaire De Dijon, Dijon, France, ³Université Bourgogne Franche Comté, LNC UMR1231, Dijon, France

Introduction: It has been suggested that dynamic arterial elastance ($E_{a_{dyn}}$) can predict decreases in arterial pressure in response to changing norepinephrine levels. The objective of this study was to determine whether $E_{a_{dyn}}$ is correlated with determinants of the vascular waterfall [critical closing pressure (CCP) and systemic arterial resistance (SARi)] in patients treated with norepinephrine.

Materials and Methods: Patients treated with norepinephrine for vasoplegia following cardiac surgery were studied. Vascular and flow parameters were recorded immediately before the norepinephrine infusion and then again once hemodynamic parameters had been stable for 15 min. The primary outcomes were $E_{a_{dyn}}$ and its associations with CCP and SARi. The secondary outcomes were the associations between $E_{a_{dyn}}$ and vascular/flow parameters.

Results: At baseline, all patients were hypotensive with $E_{a_{dyn}}$ of 0.93 [0.47;1.27]. Norepinephrine increased the arterial blood pressure, cardiac index, CCP, total peripheral resistance (TPRi), arterial elastance, and ventricular elastance and decreased $E_{a_{dyn}}$ [0.40 (0.30;0.60)] and SARi. $E_{a_{dyn}}$ was significantly associated with arterial compliance (C_A), CCP, and TPRi ($p < 0.05$).

Conclusion: In patients with vasoplegic syndrome, $E_{a_{dyn}}$ was correlated with determinants of the vascular waterfall. $E_{a_{dyn}}$ is an easy-to-read functional index of arterial load that can be used to assess the patient's macro/microcirculatory status.

Clinical Trial Registration: ClinicalTrials.gov #NCT03478709.

Keywords: dynamic arterial elastance, norepinephrine, waterfall phenomenon, vascular resistance, cardiac output

INTRODUCTION

Arterial hypotension in relation to vasoplegic syndrome is one of the most important causes of acute circulatory failure in the intensive care unit (ICU; Levin et al., 2009; Fischer and Levin, 2010). This condition is usually treated by the intravenous (IV) administration of a vasoactive agent, such as norepinephrine. The assessment of vascular tone is difficult to carry out in daily practice because the vascular parameters used (peripheral resistance and arterial compliance) did not take into account the complexity of the cardiovascular system which includes two pressure systems with a waterfall phenomenon at the microcirculatory level (Maas et al., 2012). It has been suggested that dynamic arterial elastance ($E_{a_{dyn}}$) is a marker of vascular tone that can help the physician to manage the treatment of acute vasoplegic circulatory failure (Monge Garcia et al., 2011; Pinsky, 2011; Bouchacourt et al., 2013; Cecconi et al., 2015; Guinot et al., 2015). $E_{a_{dyn}}$ is calculated using the ratio of respiratory pulse pressure variation (PPV) over the respiratory stroke volume variation (SVV). Indeed, it has been shown that $E_{a_{dyn}}$ can predict (i) the increase in mean arterial pressure (MAP) following a fluid challenge (Monge Garcia et al., 2011) and (ii) the decrease in blood pressure following a decrease in the norepinephrine dose (Guinot et al., 2015, 2017; Nguyen et al., 2021). It is a simple marker, is easy to read, and does not require additional monitoring than that carried out routinely in intensive care (Nguyen et al., 2021). However, the physiological significance of $E_{a_{dyn}}$ has never been unambiguously demonstrated. Some studies have evidenced an association between $E_{a_{dyn}}$ and several components of blood pressure: MAP, effective arterial elastance (E_A), and arterial compliance (C_A ; Guinot et al., 2017; Monge Garcia et al., 2017; Bar et al., 2018a).

The concept of vascular waterfall introduced the existence of a pressure gradient between the critical closing pressure (CCP) at the arteriolar level and the mean systemic filling pressure (MSP) at the venular level. The CCP has been described on the microcirculatory level and corresponds to the arterial pressure below which blood flow is stopped by occlusion of the arterioles (Maas et al., 2012). The CCP and indexed systemic arterial resistance (SARi) may also be markers of the vascular tone at the junction between the microcirculatory and macrocirculatory systems. It has been shown that the changes in the vascular waterfall induced by sympathetic stimulation and arterial vasodilation are similar to those observed for $E_{a_{dyn}}$ (Gaskell and Krisman, 1958; Sanders et al., 1988). The association between $E_{a_{dyn}}$ and vascular waterfall components (CCP and SARi) in vasoplegic patients in the ICU has not previously been evaluated. Given that vasomotor manipulation changes CCP and SARi, we hypothesized that $E_{a_{dyn}}$ indirectly reflects the status of these vascular waterfall components.

The main objective of the present study was to determine whether $E_{a_{dyn}}$ is associated with the vascular waterfall components in post-operative cardiac surgery patients treated with norepinephrine. The secondary objectives were to assess the putative associations between $E_{a_{dyn}}$ and several arterial load and blood flow variables.

MATERIALS AND METHODS

Ethics

The study was approved by the local independent ethics committee (Comité de Protection des Personnes Est II, Besançon, France; reference: 17/07, June 19, 2017) and was registered at Clinicaltrials.gov (NCT03478709). The present report was drafted in line with the STARD statement (Bossuyt et al., 2015). The study complied with the tenets of the Declaration of Helsinki. All patients provided their written informed consent to participation.

Patients

We performed a bicentric, prospective, observational study in the cardiothoracic and vascular ICU in two university hospitals. The inclusion criteria were as follows: adult patients admitted to the ICU and for whom the physician had decided, in accordance with the unit's care protocols, to treat post-operative vasoplegic syndrome; monitoring with a central venous access and an invasive blood pressure measurement; and the presence of a sinus heart rate (HR). Vasoplegic syndrome was defined by persistent arterial hypotension despite fluid resuscitation (Abou-Arab et al., 2018, 2020). The non-inclusion criteria were the presence of a pacemaker, atrial fibrillation, ventricular tachycardia, or ventricular fibrillation; age under 18 years; prior use of vasoactive/inotropic support; or right heart failure.

Hemodynamic Measurements

We measured the systolic blood pressure (SAP), diastolic blood pressure (DAP), and MAP using an invasive arterial catheter.

Echocardiography (using a CX50 ultrasound system and S5-1 Sector Array Ultrasound Probe Transducer, Philips Medical Systems*, Suresnes, France) was performed by a physician who was blinded to the study outcomes. The left ventricular ejection fraction (LVEF) was measured using Simpson's biplane method with a four-chamber view. The indexed stroke volume (SV_i; ml) was calculated by multiplying the left ventricular outflow tract's (LVOT's) velocity time integral (VTI) by the aortic area and dividing by body surface area (BSA). The cardiac index (CI) was calculated as SV_i × HR.

The respiratory SVV was calculated as $(SV_{max} - SV_{min}) / [(SV_{max} + SV_{min})/2] \times 100$.

Mean echocardiographic parameters were calculated from five individual beat measurements (regardless of the respiratory cycle) and analyzed retrospectively by a senior cardiologist with echocardiography certification.

MSP and CCP

The MSP and CCP were measured using the inspiratory breath-hold maneuver described by Maas et al. (2012). The measured venous and arterial pressures were plotted against cardiac output (CO), and a linear regression line was fitted to the data points. The MSP was determined by extrapolation of the central venous pressure (CVP) to zero flow on the venous return curve. The CCP was determined by extrapolation of the arterial pressure to zero flow on the ventricular output curve. Indexed SARi

(mmHg ml⁻¹.m⁻²) was calculated as (MAP – CCP)/CI. Indexed systemic venous resistance (VRi, mmHg ml⁻¹.m⁻²) was calculated as (MSP – CVP)/CI.

Left Ventricular End-Systolic Elastance, Arterial Elastance, and Ventricular-Arterial Coupling

Left ventricular end-systolic elastance (E_V) was estimated at the bedside by using the equation $E_V = \text{MAP}/(\text{ESVi}-4; \text{mmHg ml}^{-1}.\text{m}^{-2}; \text{Tanaka et al., 1998})$. This measure non-invasively used echocardiographic measurements of end-systolic volume index (ESVi). This method was used in cardiologic studies to assess the effect of vasoactive medication on ventriculo-arterial coupling (Asanoi et al., 1989; Shmuel et al., 2012).

E_A is an index that integrates the main components of arterial load, i.e., total peripheral resistance (TPRi), total net C_A , characteristic impedance, and systolic and diastolic time intervals (Tanaka et al., 1998; Ky et al., 2013). E_A was estimated by using the equation $E_A = \text{MAP}/\text{SVi}$ (mmHg ml⁻¹.m⁻²). This ratio has been shown to be a robust E_A surrogate over a wide range of hemodynamic conditions and is interchangeable in any peripheral artery (Monge Garcia et al., 2018).

Ventricular-arterial coupling was calculated as E_A/E_V ratio.

TPRi was calculated as (MAP – CVP)/CI (mmHg ml⁻¹.m⁻²), and C_A (ml mmHg⁻¹.m⁻²) was calculated as SVi/arterial pulse pressure (PP).

$E_{a_{\text{dyn}}}$ was calculated as the ratio between the PPV and the SVV (Guinot et al., 2017).

Study Protocol

The indication for treatment with norepinephrine was left to the discretion of the physician in charge of the patient. The following clinical parameters were recorded: age, gender, weight, height, body mass index, Euroscore 2, medical history, operating time, and cardiopulmonary bypass duration. After an equilibration period, baseline measurements of HR, SAP, MAP, DAP, MSP, TPRi, SVi, CI, SVV, PPV, and vascular determinants of waterfall were recorded. A second (post-norepinephrine) set of measurements was recorded after 15 min of hemodynamic stability (defined as a change in MAP of less than 10%). All patients had been sedated by continuous infusion of propofol and were fully adapted to mechanical ventilation. All patients were mechanically ventilated in pressure-controlled mode with a positive end-expiratory pressure (PEEP) of 5 cmH₂O. Ventilator settings and the propofol infusion rate were the same for the two sets of measurements.

Statistical Methods

We calculated that a sample size of 14 patients would allow to demonstrate a correlation between $E_{a_{\text{dyn}}}$ and CCP higher than 0.5, using Spearman's correlation coefficient with a risk alpha of 5% and a power of 80%. The distribution of the variables was assessed using the D'Agostino-Pearson test. Data are expressed as proportion (%), mean (standard deviation), or median [IQR] (interquartile range) as appropriate. Paired data were compared using Wilcoxon signed rank test. Linear correlations were tested using Spearman's rank method.

Associations between arterial load and blood flow parameters measurement considering both time points were assessed using mixed linear modeling. Only bivariate models were carried out. $E_{a_{\text{dyn}}}$ was the dependent variable; each hemodynamic variable was used as fixed effect, and individual was used as random intercept. The normality of the distribution of random effects and of the model residual were graphically checked. Because this analysis was exploratory, slight disparities between those distributions and normality were tolerated. The threshold for statistical significance was set to 0.05. Statistical analysis was performed using RStudio Version 1.1.447-2009-2018 RStudio, Inc., and SPSS® software (version 24, IBM, New York, NY, United States).

RESULTS

Twenty patients were included during the study period. Three were excluded because of incomplete data, and so 17 were included in the final analysis. Most of the study participants were male (80%). The median age was 68 [50;71], and the median Euroscore 2 was 2.3 [1.5;4.3] (Table 1). The median dose of norepinephrine was 0.08 gamma kg⁻¹.min⁻¹ [0.06;0.11]. None of the patients had been treated with inotropic agents or vasopressors before enrolment in the study.

At baseline, patients had arterial hypotension, with low CCP and CI values, and high SARI, SVV, PPV, and $E_{a_{\text{dyn}}}$ values (Table 2). Infusion of norepinephrine significantly increased SAP, DAP, MAP, PP, CVP, CCP, TPRi, VRi, SVV, SVi, CI, E_A , and E_V and significantly decreased $E_{a_{\text{dyn}}}$, PPV, C_A , and SARI (Table 2). These changes did not differ between non-smoker and smoker patients ($p = 0.45$).

TABLE 1 | Demographic data.

Variables	(N = 20)
Age (years)	68 [50–71]
Male (%)	16 (80%)
Weight (kg)	85 ± 18
Height (cm)	175 ± 9
BMI (kg·m ⁻²)	27.9 ± 4.8
Euroscore 2 (%)	2.3 [1.5–4.3]
Medical history, n (%)	
High blood pressure	13 (65%)
Diabetes	0 (0%)
Coronary artery disease	5 (25%)
Smoking	11 (55%)
Dyslipidemia	6 (30%)
Surgery, n (%)	
Valve surgery	10 (50%)
Coronary artery bypass graft	7 (35%)
Both of the above procedures	3 (15%)
Cardiopulmonary bypass duration (min)	117 [91–158]
Length of stay in the ICU (days)	3 [2–5]
Length of stay in hospital (days)	11 ± 3

The data were quoted as the number (%), mean ± standard deviation SD, or the median [interquartile range, IQR], as appropriate. BMI, body mass index; ICU, intensive care unit.

TABLE 2 | Comparison of hemodynamic parameters at baseline and after the norepinephrine injection.

Hemodynamic variables	Baseline	Post-injection	<i>p</i>
HR (bpm)	75.5 [63.0;91.2]	79.0 [64.2;91.8]	0.491
SAP (mmHg)	89 [84; 94]	122 [114;130]	0.001
DAP (mmHg)	46.5 [41.0;50.2]	62.5 [55.5;67.2]	0.001
MAP (mmHg)	60.5 [56.0;66.0]	83.0 [76.5;89.0]	0.001
PP (mmHg)	41.5 [35.0;46.5]	57.5 [49.8;68.2]	<0.001
CVP (mmHg)	7.00 [5.50;9.00]	8.50 [5.75;9.25]	<0.001
SVV (%)	17.2 [15.0;23.7]	25.0 [21.9;33.3]	0.002
PPV (%)	14.5 [10.5;18.2]	9.50 [7.00;14.5]	0.027
Ea _{dyn}	0.93 [0.47;1.27]	0.40 [0.30;0.60]	0.042
MSP (mmHg)	23.0 [18.5;26.0]	19.0 [14.8;24.0]	0.149
VRi (mmHg ml ⁻¹ .m ⁻²)	2.79 [2.63;2.98]	3.06 [2.76;3.26]	0.005
CCP (mmHg)	26.1 [19.4;29.1]	54.8 [39.8;70.3]	0.001
SARi (mmHg ml ⁻¹ .m ⁻²)	18.4 [14.5;25.8]	11.4 [6.06;18.5]	0.002
CCP-MSP (mmHg)	5.74 [-3.8;11.1]	42.0 [17.1;48.0]	0.001
SVi (ml.m ⁻²)	23.0 [17.5;28.8]	28.0 [19.0;36.0]	0.034
CI (ml min ⁻¹ .m ⁻²)	1.66 [1.56;2.09]	1.96 [1.82;2.32]	0.010
E _A (mmHg ml ⁻¹ .m ⁻²)	1.66 [1.41;2.31]	1.75 [1.50;3.25]	0.003
Total C _A (ml mmHg ⁻¹ .m ⁻²)	1.27 [1.01;1.61]	0.95 [0.78;1.22]	0.016
TPRi (mmHg ml ⁻¹ .m ⁻²)	29.0 [24.5;39.0]	36.0 [29.2;43.0]	0.004
Ventricular elastance (E _v ; mmHg ml ⁻¹ .m ⁻²)	1.2 [0.9;1.9]	1.74 [1.1;2.4]	0.002
E _A /E _v ratio (units)	1.4 [1.1;1.9]	1.3 [1;1.9]	0.215

Values are expressed as the mean ± SD or the median [IQR]. C_A, arterial compliance; CI, cardiac index; CVP, central venous pressure; DAP, diastolic arterial pressure; E_A, arterial elastance; Ea_{dyn}, dynamic arterial elastance; E_v, ventricular elastance; PPV, pulse pressure variation; HR, heart rate; MAP, mean arterial pressure; CCP, critical closing pressure; PP, pulse pressure; TPRi, indexed total peripheral resistance; SAP, systolic arterial pressure; SARi, indexed systemic arterial resistance; SVi, indexed stroke volume; SVV, stroke volume variation; VRi, indexed systemic venous resistance. The value of *p* refers to the within-group (before/after) comparison. Bold values are values with *p* < 0.05.

In the overall study population, Ea_{dyn} was significantly and negatively correlated with CCP ($r = -0.557$, $p = 0.0017$), CCP-MSP ($r = -0.515$, $p = 0.003$), SARi ($r = -0.487$, $p = 0.003$), SAP ($r = -0.402$, $p = 0.004$), DAP ($r = -0.484$, $p = 0.005$), MAP ($r = -0.498$, $p = 0.004$), E_A ($r = -0.347$, $p = 0.05$), C_A ($r = -0.406$, $p = 0.021$), and TPRi ($r = -0.396$, $p = 0.025$; **Figures 1, 2**). Ea_{dyn} was not correlated with LVEF ($r = 0.012$, $p = 0.945$).

The linear mixed model used to analyze the contribution of arterial and blood flow parameters to Ea_{dyn} demonstrated that Ea_{dyn} was significantly associated with C_A, blood pressure, CCP, and TPRi (**Table 3**).

DISCUSSION

The present study's main finding was the association between Ea_{dyn} and determinants of the vascular waterfall (CCP and SARi). Furthermore, we confirmed that Ea_{dyn} is associated with several components of arterial load (C_A and TPRi) and blood pressure (MAP and CCP).



According to the concept of macrocirculatory-microcirculatory coherence, any change in the macrocirculation will affect the microcirculation (Bennett et al., 2018). Hence, resuscitation procedures intended to correct macrocirculatory hemodynamic variables will also modify regional and microcirculatory

perfusion and oxygen delivery to the tissues (Ince, 2015). The CCP was described at the microcirculatory level and represents the arterial pressure below which blood flow is stopped by occlusion of the arterioles (Versluis et al., 2001; Maas et al., 2012). The vascular waterfall is thought to maintain tissue perfusion when blood flow is abnormally low. The CCP values observed in the present study were similar to those published in the literature (Jellinek et al., 2000; Schipke et al., 2003; Kottenberg-Assenmacher et al., 2009). As expected, norepinephrine increased CCP, the vascular waterfall (i.e., the CCP-MSP difference), and thus the level of tissue perfusion.

When arterial hypotension occurs, the main adaptive mechanism is the maintenance of blood flow to the tissues. In this respect, it has been suggested that several adaptive hemodynamic factors are activated so as to maintain the blood flow redistribution (De Hert, 2012). The blood pressure increase with norepinephrine treatment might restore the blood pressure and thus diminish the contribution of these hemodynamic factors to changes in tissue perfusion. Hence, we observed a decrease in SARi, which might be due to the redistribution of blood flow to non-perfused capillaries (microvascular recruitment) with norepinephrine, and a decrease in the adaptive phenomena described above (Georger et al., 2010; De Backer et al., 2014).

The changes in Ea_{dyn} after norepinephrine infusion contrast with those observed upon norepinephrine weaning in the patients with sepsis or after cardiac surgery (Guinot et al., 2017). Likewise, studies of Ea_{dyn} in the operating theater and in the ICU have given conflicting results (Guinot et al., 2015; Lanchon et al., 2017; Bar et al., 2018b). These disparities may be due to several factors, e.g., the non-linearity of the blood pressure-volume relationship (Sanders et al., 1988), the cardiovascular effects of norepinephrine (as an alpha and beta-agonist; Guinot et al., 2018) and the underlying disease (vasoplegia is associated with altered autonomic function; Bernardelli et al., 2016), and thus changes in the interactions between the arterial load and blood pressure. Hence, Ea_{dyn} decreased with norepinephrine infusion probably because of a decrease in C_A, an increase in left ventricular inotropy (E_v), and a decrease in arteriolar vasomotor tone (SARi; **Figure 3**).

As demonstrated by the literature data, C_A is probably the main contributor to Ea_{dyn} (Monge García et al., 2017; Bar et al., 2018a). It has been shown that norepinephrine infusion can restore C_A (Nguyen et al., 2018). Fluid expansion does not change C_A in patients treated with norepinephrine but decreases C_A in patients not treated with norepinephrine. Likewise, norepinephrine weaning is not associated with a change of C_A in non-pressure-responder patients (Bar et al., 2018b). Thus, in norepinephrine-treated patients, changes in the arterial load may mainly be driven by the resistive component (i.e., SARi and TPRi; Bernardelli et al., 2016; Nguyen et al., 2018). One can suppose that in patients treated with vasopressor, Ea_{dyn} mainly reflects vascular resistance because C_A is fixed. This hypothesis may also explain the ability of Ea_{dyn} (when measured with an uncalibrated CO device) to predict arterial pressure changes (Bar et al., 2018b). The arterial load and Ea_{dyn} may vary in a complex manner as a function of the



therapeutic intervention while decreasing Ea_{dyn} 's ability to predict blood pressure changes. The main negative studies concerning Ea_{dyn} were performed in the operating theater, and the positive studies were performed in the ICU (i.e., where most of the patients had received vasopressors; Vos et al., 2013;


Lanchon et al., 2017; Guarracino et al., 2019). In practice, Ea_{dyn} could be considered as an easy-to-read indicator that can help the physician to choose the optimal hemodynamic strategy at the bedside. This index could be used to understand the interaction between cardiac function and arterial load, the effects of hemodynamic treatment on components of arterial load, coherence between macrocirculation and microcirculation (i.e., the vascular waterfall's status), and thus to determine which treatments could be used or withdrawn. Guarracino et al. (2019) have demonstrated that Ea_{dyn} predicts changes in MAP upon fluid expansion or vasopressor administration. In contrast to the complex procedures required to measure the vascular waterfall and arterial load variables, Ea_{dyn} is easy to measure with hemodynamic devices – even an uncalibrated pulse contour analysis device (Bar et al., 2018b).

TABLE 3 | Estimated values of fixed effects (arterial load and cardiac parameters) on Ea_{dyn} according to a linear mixed-effects model analysis.

Covariates	Ea_{dyn}	<i>p</i>
Arterial covariates		
SAP (mmHg)	-0.02 (0.01)	0.040
DAP (mmHg)	-0.03 (0.01)	0.040
MAP (mmHg)	-0.03 (0.01)	0.030
PP (mmHg)	0.19 (0.04)	0.190
CCP (mmHg)	-0.55 (0.02)	0.020
SARi (mmHg ml ⁻¹ .m ⁻²)	-0.11 (0.04)	0.730
E_A (mmHg ml ⁻¹ .m ⁻²)	-0.43 (0.22)	0.070
C_A (ml mmHg ⁻¹ .m ⁻²)	1.25 (0.33)	0.010
TPRi (mmHg ml ⁻¹ .m ⁻²)	-0.05 (0.02)	0.020
Norepinephrine dose (g kg ⁻¹ min ⁻¹)	-5.9 (3.3)	0.090
Cardiac covariates		
HR (beats min ⁻¹)	-0.01 (0.01)	0.230
LVEF (%)	0.016 (0.011)	0.670
E_V (mmHg ml ⁻¹ .m ⁻²)	-0.17 (0.22)	0.330
E_A/E_V (units)	-0.10 (0.038)	0.610
CI (l min ⁻¹ .m ⁻²)	0.89 (0.36)	0.020

Data are presented as an estimated value (SE). C_A , arterial compliance; CI, cardiac index; CVP, central venous pressure; DAP, diastolic arterial pressure; E_A , arterial elastance; Ea_{dyn} , dynamic arterial elastance; E_V , ventricular elastance; NE, norepinephrine; HR, heart rate; LVEF, left ventricular ejection fraction; MAP, mean arterial pressure; MSP, mean systemic pressure; CCP, critical closing pressure; PP, pulse pressure; TPRi, indexed total peripheral resistance; SAP, systolic arterial pressure; SARi, indexed systemic arterial resistance. Bold values are values with $p < 0.05$.

The present study had several limitations. Even though we predetermined the sample size, the present results included a small number of patients and will require external validation in another cohort. Our results were obtained in a group of patients with arterial hypotension treated with norepinephrine and must now be confirmed in patients treated with fluid expansion or inotropes. We did not include a control group, i.e., using IV fluids because fluid infusion may be unable to treat vasoplegia, and it seems difficult to compare the effects of fluid on vasomotor tone and those of norepinephrine in two populations who do not suffer the same hemodynamic underlying disease. To avoid mathematical coupling, we specifically chose to calculate Ea_{dyn} from two different signals. The arterial load assessment was based on a two-element Windkessel model and integrative simplification (Westerhof et al., 2009). More precise models (include arterial impedance and wave reflection) have



since been developed. However, the performance of these measurements at the bedside would be challenging. The E_A calculated from MAP was demonstrated to be a robust surrogate over a wide range of hemodynamic conditions and can be applied to any peripheral artery (Monge Garcia et al., 2018). The method that we used to calculate E_V and E_A can be criticized because we did not use a high-fidelity catheter for ventricular pressure measurement; however, our approach has been studied previously (Huette et al., 2020; Nguyen et al., 2020). Likewise, we used a standardized inspiratory breath-hold maneuver. This method has been validated in ICU patients, and the values obtained in the present study were similar to those determined with other methods (Maas et al., 2012). Finally, there was heterogeneity in the study participants because their cause for surgery was not the same, but all patients suffer the same hemodynamic disease, and the inclusion criteria allow to homogenize the studied population.

In hypotensive patients treated with norepinephrine, $E_{a_{dyn}}$ may indirectly reflect the vascular waterfall phenomenon and changes in one or both of its determinants (CCP and SARI) following treatment. Hence, one can consider that $E_{a_{dyn}}$ reflects vasomotor tone in hypotensive patients treated with norepinephrine. In view of our present results and the literature data, one can further consider that $E_{a_{dyn}}$ indirectly

reflects macro-microcirculatory coherence during acute circulatory failure.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by Comité de Protection des Personnes Est II, Besançon, France; reference: 17/07, June 19th, 2017. The patients/participants provided their written informed consent to participate in this study.

AUTHOR CONTRIBUTIONS

SB, MN, P-GG, OA-A, and BB designed the study and interpreted the data. SB and P-GG analyzed the data. SB, P-GG, and HD drafted the manuscript for important intellectual content. All authors contributed to the article and approved the submitted version.

REFERENCES

- Abou-Arab, O., Kamel, S., Beyls, C., Huette, P., Bar, S., Lorne, E., et al. (2020). Vasoplegia after cardiac surgery is associated with endothelial glyocalyx alterations. *J. Cardiothorac. Vasc. Anesth.* 34, 900–905. doi: 10.1053/j.jvca.2019.09.004
- Abou-Arab, O., Martineau, L., Bar, S., Huette, P., Amar, A. B., Caus, T., et al. (2018). Postoperative vasoplegic syndrome is associated with impaired endothelial vasomotor response in cardiac surgery: a prospective, observational study. *J. Cardiothorac. Vasc. Anesth.* 32, 2218–2224. doi: 10.1053/j.jvca.2018.02.006
- Asanoi, H., Sasayama, S., and Kameyama, T. (1989). Ventriculoarterial coupling in normal and failing heart in humans. *Circ. Res.* 65, 483–493. doi: 10.1161/01.RES.65.2.483
- Bar, S., Huette, P., Abou-Arab, O., Dupont, H., Lorne, E., and Guinot, P.-G. (2018a). Dynamic arterial elastance might not be an indicator of ventriculoarterial coupling. Comment on Br J Anaesth 2017; 118: 938–46. *Br. J. Anaesth.* 121, 672–673. doi: 10.1016/j.bja.2018.03.027
- Bar, S., Levie, F., Abou Arab, O., Badoux, L., Mahjoub, Y., Dupont, H., et al. (2018b). Dynamic arterial elastance measured by uncalibrated pulse contour analysis predicts arterial-pressure response to a decrease in norepinephrine. *Br. J. Anaesth.* 121, 534–540. doi: 10.1016/j.bja.2018.01.032
- Bennett, V. A., Vidouris, A., and Cecconi, M. (2018). Effects of fluids on the macro- and microcirculations. *Crit. Care* 22:74. doi: 10.1186/s13054-018-1993-1
- Bernardelli, A. K., Da Silva, R. d. C. V. d. A. F., Corrêa, T., and Da Silva-Santos, J. E. (2016). Vasoplegia in sepsis depends on the vascular system, vasopressor, and time-point: a comparative evaluation in vessels from rats subjected to the cecal ligation puncture model. *Can. J. Physiol. Pharmacol.* 94, 1227–1236. doi: 10.1139/cjpp-2015-0514
- Bossuyt, P. M., Reitsma, J. B., Bruns, D. E., Gatsonis, C. A., Glasziou, P. P., Irwig, L., et al. (2015). STARD 2015: an updated list of essential items for reporting diagnostic accuracy studies. *BMJ* 351, 1–9. doi: 10.1136/bmj.h5527
- Bouchacourt, J. P., Riva, J. A., and Grignola, J. C. (2013). The increase of vasomotor tone avoids the ability of the dynamic preload indicators to estimate fluid responsiveness. *BMC Anesthesiol.* 13:41. doi: 10.1186/1471-2253-13-41
- Cecconi, M., García, M. I. M., Romero, M. G., Mellinshoff, J., Caliendo, F., Grounds, R. M., et al. (2015). The use of pulse pressure variation and stroke volume variation in spontaneously breathing patients to assess dynamic arterial elastance and to predict arterial pressure response to fluid administration. *Anesth. Analg.* 120, 76–84. doi: 10.1213/ANE.0000000000000442
- De Backer, D., Orbeago Cortes, D., Donadello, K., and Vincent, J.-L. (2014). Pathophysiology of microcirculatory dysfunction and the pathogenesis of septic shock. *Virulence* 5, 73–79. doi: 10.4161/viru.26482
- De Hert, S. (2012). Physiology of hemodynamic homeostasis. *Best Pract. Res. Clin. Anaesthesiol.* 26, 409–419. doi: 10.1016/j.bpa.2012.10.004
- Fischer, G. W., and Levin, M. A. (2010). Vasoplegia during cardiac surgery: current concepts and management. *Semin. Thorac. Cardiovasc. Surg.* 22, 140–144. doi: 10.1053/j.semtcvs.2010.09.007
- Gaskell, P., and Krisman, A. M. (1958). Critical closing pressure of vessels supplying the capillary loops of the nailfold. *Circ. Res.* 6, 461–467. doi: 10.1161/01.RES.6.4.461
- Georger, J.-F., Hamzaoui, O., Chaari, A., Maizel, J., Richard, C., and Teboul, J.-L. (2010). Restoring arterial pressure with norepinephrine improves muscle tissue oxygenation assessed by near-infrared spectroscopy in severely hypotensive septic patients. *Intensive Care Med.* 36, 1882–1889. doi: 10.1007/s00134-010-2013-3
- Guarracino, F., Bertini, P., and Pinsky, M. R. (2019). Cardiovascular determinants of resuscitation from sepsis and septic shock. *Crit. Care* 23:118. doi: 10.1186/s13054-019-2414-9
- Guinot, P.-G., Abou-Arab, O., Guilbart, M., Bar, S., Zogheib, E., Daher, M., et al. (2017). Monitoring dynamic arterial elastance as a means of decreasing the duration of norepinephrine treatment in vasoplegic syndrome following cardiac surgery: a prospective, randomized trial. *Intensive Care Med.* 43, 643–651. doi: 10.1007/s00134-016-4666-z
- Guinot, P.-G., Bernard, E., Levrard, M., Dupont, H., and Lorne, E. (2015). Dynamic arterial elastance predicts mean arterial pressure decrease associated with decreasing norepinephrine dosage in septic shock. *Crit. Care* 19:14. doi: 10.1186/s13054-014-0732-5
- Guinot, P.-G., Longrois, D., Kamel, S., Lorne, E., and Dupont, H. (2018). Ventriculoarterial coupling analysis predicts the hemodynamic response to norepinephrine in hypotensive postoperative patients: a prospective observational study. *Crit. Care Med.* 46, e17–e25. doi: 10.1097/CCM.0000000000002772

- Huette, P., Abou-Arab, O., Longrois, D., and Guinot, P.-G. (2020). Fluid expansion improve ventriculo-arterial coupling in preload-dependent patients: a prospective observational study. *BMC Anesthesiol.* 20:171. doi: 10.1186/s12871-020-01087-7
- Ince, C. (2015). Hemodynamic coherence and the rationale for monitoring the microcirculation. *Crit. Care* 19, 1–13. doi: 10.1186/cc14726
- Jellinek, H., Krenn, H., Oczenski, W., Veit, F., Schwarz, S., and Fitzgerald, R. D. (2000). Influence of positive airway pressure on the pressure gradient for venous return in humans. *J. Appl. Physiol.* 88, 926–932. doi: 10.1152/jappl.2000.88.3.926
- Kottenberg-Assenmacher, E., Aleksic, I., Eckholt, M., Lehmann, N., and Peters, J. (2009). Critical closing pressure as the arterial downstream pressure with the heart beating and during circulatory arrest. *Anesthesiology* 110, 370–379. doi: 10.1097/ALN.0b013e3181942c99
- Ky, B., French, B., May Khan, A., Plappert, T., Wang, A., Chirinos, J. A., et al. (2013). Ventricular-arterial coupling, remodeling, and prognosis in chronic heart failure. *J. Am. Coll. Cardiol.* 62, 1165–1172. doi: 10.1016/j.jacc.2013.03.085
- Lancho, R., Nouette-Gaulain, K., Stecken, L., Sesay, M., Lefrant, J.-Y., and Biais, M. (2017). Dynamic arterial elastance obtained using arterial signal does not predict an increase in arterial pressure after a volume expansion in the operating room. *Anaesthesia Critical Care & Pain Medicine* 36, 377–382. doi: 10.1016/j.accpm.2017.05.001
- Levin, M. A., Lin, H.-M., Castillo, J. G., Adams, D. H., Reich, D. L., and Fischer, G. W. (2009). Early on-cardiopulmonary bypass hypotension and other factors associated with vasoplegic syndrome. *Circulation* 120, 1664–1671. doi: 10.1161/CIRCULATIONAHA.108.814533
- Maas, J. J., de Wilde, R. B., Aarts, L. P., Pinsky, M. R., and Jansen, J. R. (2012). Determination of vascular waterfall phenomenon by bedside measurement of mean systemic filling pressure and critical closing pressure in the intensive care unit. *Anesth. Analg.* 114, 803–810. doi: 10.1213/ANE.0b013e318247fa44
- Monge Garcia, M. I., Gil Cano, A., and Gracia Romero, M. (2011). Dynamic arterial elastance to predict arterial pressure response to volume loading in preload-dependent patients. *Crit. Care* 15:R15. doi: 10.1186/cc9420
- Monge García, M. I., Guijo González, P., Gracia Romero, M., Gil Cano, A., Rhodes, A., Grounds, R. M., et al. (2017). Effects of arterial load variations on dynamic arterial elastance: an experimental study. *Br. J. Anaesth.* 118, 938–946. doi: 10.1093/bja/aex070
- Monge Garcia, M. I., Jian, Z., Settels, J. J., Hatib, F., Cecconi, M., and Pinsky, M. R. (2018). Reliability of effective arterial elastance using peripheral arterial pressure as surrogate for left ventricular end-systolic pressure. *J. Clin. Monit. Comput.* 33, 803–813. doi: 10.1007/s10877-018-0236-y
- Nguyen, M., Abou-Arab, O., Bar, S., Dupont, H., Bouhemad, B., and Guinot, P.-G. (2021). Echocardiographic measure of dynamic arterial elastance predict pressure response during norepinephrine weaning: an observational study. *Sci. Rep.* 11:2853. doi: 10.1038/s41598-021-82408-9
- Nguyen, M., Berhoud, V., Bartamian, L., Martin, A., Ellouze, O., Bouhemad, B., et al. (2020). Agreement between different non-invasive methods of ventricular elastance assessment for the monitoring of ventricular-arterial coupling in intensive care. *J. Clin. Monit. Comput.* 34, 893–901. doi: 10.1007/s10877-019-00397-7
- Nguyen, M., Fischer, M.-O., Mallat, J., Marc, J., Lorne, E., Bouhemad, B., et al. (2018). Fluid responsiveness depend on arterial load and vasopressor use in ICU patients. ESICM LIVES 2018: Paris, France. 20–24 October 2018. *Intensive Care Medicine Experimental* 6:59. doi: 10.1186/s40635-018-0201-6
- Pinsky, M. R. (2011). Defining the boundaries of bedside pulse contour analysis: dynamic arterial elastance. *Crit. Care* 15:120. doi: 10.1186/cc9986
- Sanders, J. S., Ferguson, D. W., and Mark, A. L. (1988). Arterial baroreflex control of sympathetic nerve activity during elevation of blood pressure in normal man: dominance of aortic baroreflexes. *Circulation* 77, 279–288. doi: 10.1161/01.CIR.77.2.279
- Schipke, J. D., Heusch, G., Sani, A. P., Gams, E., and Winter, J. (2003). Static filling pressure in patients during induced ventricular fibrillation. *Am. J. Phys. Heart Circ. Phys.* 285, H2510–H2515. doi: 10.1152/ajpheart.00604.2003
- Shmuel, S., Redfield, M. M., From, A. M., Paul, S., Nishimura, R. A., and Borlaug, B. A. (2012). Effects of vasodilation in heart failure with preserved or reduced ejection fraction. *J. Am. Coll. Cardiol.* 59, 442–451. doi: 10.1016/j.jacc.2011.09.062
- Tanaka, K., Oshita, S., Kitahata, H., Kimura, H., Kawahito, S., Park, Y. C., et al. (1998). Effects of nicardipine on ventriculo-arterial coupling in humans. *Br. J. Anaesth.* 81, 180–185. doi: 10.1093/bja/81.2.180
- Versluis, J. P., Heslinga, J. W., Sipkema, P., and Westerhof, N. (2001). Microvascular pressure measurement reveals a coronary vascular waterfall in arterioles larger than 110 μm . *Am. J. Phys. Heart Circ. Phys.* 281, H1913–H1918. doi: 10.1152/ajpheart.2001.281.5.H1913
- Vos, J. J., Kalmar, A. F., Struys, M. M. R. F., Wietasch, J. K. G., Hendriks, H. G. D., and Scheeren, T. W. L. (2013). Comparison of arterial pressure and plethysmographic waveform-based dynamic preload variables in assessing fluid responsiveness and dynamic arterial tone in patients undergoing major hepatic resection. *Br. J. Anaesth.* 110, 940–946. doi: 10.1093/bja/aes508
- Westerhof, N., Lankhaar, J.-W., and Westerhof, B. E. (2009). The arterial Windkessel. *Med. Biol. Eng. Comput.* 47, 131–141. doi: 10.1007/s11517-008-0359-2

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Bar, Nguyen, Abou-Arab, Dupont, Bouhemad and Guinot. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

GLOSSARY

Term	Definition
C_A	Arterial compliance
CCP	Critical closing pressure
CI	Cardiac index
CO	Cardiac output
CVP	Central venous pressure
DAP	Diastolic arterial pressure
E_A	Arterial elastance
$E_{a,yn}$	Dynamic arterial elastance
E_V	Ventricular elastance
HR	Heart rate
LVEF	Left ventricular ejection fraction
MAP	Mean arterial pressure
MSP	Mean systemic pressure
PP	Pulse pressure
PPV	Pulse pressure variation
TPRi	Indexed total peripheral resistance
SAP	Systolic arterial pressure
SARi	Indexed systemic arterial resistance
SVi	Indexed stroke volume
SVV	Stroke volume variation
VRI	Indexed systemic venous resistance